

Effect of Mean Annual Changeability of Air Temperature, Surface Ozone Concentration and Galactic Cosmic Rays Intensity on the Mortality of Tbilisi City Population

**¹Avtandil G. Amiranashvili, ¹Teimuraz S. Bakradze,
²Nino T. Berianidze, ³Nino D. Japaridze, ^{3,4}Ketevan R. Khazaradze**

¹*M. Nodia Institute of Geophysics of I. Javakhishvili Tbilisi State University, e-mail:*

avtandilamiranashvili@gmail.com

²*National Science Library of Georgia*

³*Ministry of Labor, Health and Social Affairs of Georgia*

⁴*Georgian State Teaching University of Physical Education and Sport*

ABSTRACT

The results of a study of the effect of the annual changeability of air temperature, surface ozone concentration and neutron component of galactic cosmic rays intensity on the mortality of the population of Tbilisi city in 1984-2010 are presented. The statistical characteristics of the investigated time-series are studied. In particular, it was found that within the variation range the contribution of the studied parameters to mortality variability is as follows: a random component of air temperature - 8.5%, real values of surface ozone concentration and cosmic ray intensity - 20.9% and 16.5%, respectively.

Key Words: Air temperature, surface ozone concentration, galactic cosmic rays, mortality, ecology, bioclimatology, medical meteorology.

Introduction

The meteorological, biometeorological, bioclimatic, geophysical and other parameters, which sufficiently affect the human beings, are the followings.

Separate meteorological and geophysical elements, space weather parameters and its combinations: air temperature [1-10], humidity, wind speed, atmospheric pressure, solar activity (Wolf's number), the geomagnetic fields, solar radiation, the cosmic rays [11-18], light ions, aerosols, ozone, other air toxic admixtures and etc. [19-26].

Different simple thermal indices involve more than one climatological parameter and consider the combined effects (air equivalent- effective temperature EET, Equivalent temperature (TEK), Wet-bulb-globe temperature (WBGT), Tourism Climate Index (TCI) [27-32]) and others.

The indices derived from energy budget models:

Physiologically Equivalent Temperature (PET), Standard Effective Temperature (SET), Physiological Subjective Temperature and Subjective Temperature (MENEX), the Universal Thermal Climate Index (UTCI) etc. [30, 33-36]).

Generally, the human's health is primarily affected by the lifestyle (50 – 55%), then – by the environment (25 – 30%), and finally – by heritage and medical care [19]. Additional anthropogenic load on the biosphere increases the level of above mentioned risk - factors influencing on human health and life [19, 24, 25].

The effects of the action of environmental factors on human health have different scales - from minute, hour, day, decade and month to the seasonal and annual [37-39]. For example, periodicity of 7 and

3, 5 day of mortality from the cardiovascular diseases (CVD) is established in the work [37]. Results of investigation of influence of monthly average values of air Equivalent-Effective Temperature EET and is represented monthly duration of magnetic storms D on the health of the population of Tbilisi city [27]. The analysis of regression connections of mortality from the CVD with the EET and D showed that the contribution of each of the variables into changeability of mortality is the following. In the range EET from -5° to 4.6° : EET – 8.6%, D – 22.2%; in the range EET from 5.2° to 21.8° : T - 26.3%, D - is insignificant [27].

In work [19] is shown, that days situation together with air pollution by ozone in smog, the ozone forming gases and the aerosols under the conditions of Tbilisi an essential effect on human health have a variation in such factors as the thermal regime of air, atmospheric pressure, cosmic rays. Thus, increased surface ozone concentrations on the average growth of annual mortality of the inhabitants of Tbilisi city by 1680 people. This is equal to 14.1 % of entire average annual mortality of the population of Tbilisi, which is approximately 3 times higher than the same indices for the advanced countries [19, 25, 26].

In this work are presented the results of a study of the effect of the annual changeability of air temperature, surface ozone concentration and neutron component of galactic cosmic rays intensity on the mortality of the population Tbilisi city in 1984-2010.

Material and methods

In the work are used the data of National Statistics Office of Georgia about the common mortality of the population of Tbilisi city. The common annual mortality (M) of population to 1000 inhabitants is normalized [25].

The measurements of surface ozone concentration (SOC) were conducted by the electro chemical ozone instrument OMG-200. Here are presented observational data for 15 hours [40]. The unit of the ozone measurement is mcg/m^3 .

The data of the Hydro meteorological Service of Georgia about the annual values of air temperature (T, $^{\circ}\text{C}$) in Tbilisi are used. Information about annual values of intensity of neutron component of galactic cosmic rays (CR, impulse/hour) is obtained at the Cosmic Rays Observatory of M. Nodia Institute of Geophysics. The observation period is 1984-2100.

In the proposed work the analysis of data is carried out with the use of the standard statistical analysis methods of random events and mathematical statistic methods for the non accidental time-series of observations [41, 42].

The following designations will be used below: Min – minimal values, Max - maximal values, Range - variational scope, St Dev- standard deviation, R - coefficient of linear correlation, R^2 –coefficient of determination, K_{DW} – Durbin-Watson statistic, α - the level of significance. As in [43], Res – residual component, Real - measured data. The curve of trend is equation of the regression of the connection of the investigated parameter with the time at the significant value of the determination coefficient and such values of K_{DW} , where the residual values are accidental.

Background component is usually entered into the curve of trend. The value of background component is most frequently unknown. From the physical considerations, random component can be represented in the form: $\text{Rand} = \text{Res} + \text{absolute value of the min value of Res}$. In this case random components have positive values with the minimum value = 0 (if there would be known the value of background component, that min Rand will be = Back). Accordingly, Trend + Back (sum of the trend and background components of time series) will be curve of equation of the regression of the connection of the investigated parameter with the time minus absolute value of the min value of Res. So, $\text{Real} = (\text{Trend} + \text{Back}) + \text{Res}$.

Below in the text the dimensionality of the investigated parameters are omitted

Results of detailed statistical analysis of the changeability of mean annual values of SOC in Tbilisi in 1984-2010 in [25, 40] were represented. In particular, the changeability of the indicated time series is described by the fourth power polynomial. An increase in the SOC in the period from 1984 through 1995-1997 was observed, then - decrease. Thus, in average: in 1984 $\text{SOC} = 37 \text{ mcg}/\text{m}^3$, into 1998 – $58 \text{ mcg}/\text{m}^3$, into 2010 – $40 \text{ mcg}/\text{m}^3$. There was presented [25] Information about changeability of M in the investigation period of time in Tbilisi in.

This work is the continuation of work [25]. In it, together with the action of surface ozone (both the direct action and the indicator of the pollution of atmosphere) on the mortality of the population of Tbilisi, the role of variations in the air temperature and is examined intensity of galactic cosmic rays in the changeability of values M.

Results and discussion

The results in tables 1-4 and fig. 1-2 are given.

Table 1

Characteristics of trend of M, SOC, T and CR in Tbilisi in 1984-2010

Variable	Form of the equation of regression	R ² (with year)	K _{DW}
M	Fifth power polynomial [25]	0.86 ($\alpha=0.0001$)	1.45 ($\alpha=0.05$)
SOC	Fourth power polynomial [25, 40]	0.82 ($\alpha=0.0001$)	2.03 ($\alpha=0.05$)
T	Linear	0.13 ($\alpha=0.06$)	1.55 ($\alpha=0.05$)
CR	Seventh power polynomial	0.82 ($\alpha=0.0001$)	1.89 ($\alpha=0.05$)

In table 1, is presented the data about characteristics of trend of mortality, surface ozone concentration, air temperature and cosmic rays intensity in Tbilisi in 1984-2010. As follows from table 1 trends of M, SOC, and CR take the form of fifth power, fourth power and seventh power polynomials; trend of T is linear (corresponding values of R² and K_{DW}).




Fig. 1. Trend of the average annual air temperature in Tbilisi in 1984-2010.




Fig. 2. Trend of the average annual intensity of galactic cosmic rays in Tbilisi in 1984-2010.

For the clarity in fig. 1 and 2 are presented the curves of real data, (trend+ background) and random components of time-series of mean annual air temperature and intensity of galactic cosmic rays in Tbilisi in

1984-2010. Analogous data for M and SOC in [25, 40] are given.

It should be noted that the trend of the average annual air temperature in Tbilisi into different time intervals had different nature. So, in 1963-1990 [43], 1878-1997 [44], 1906-1995 [45], 1907-2006 [46], 1957-2006 [47] the trend of T was linear. In the more prolonged period of time (1850-2012 [48]) the trend of T in Tbilisi took the form of the fourth power polynomial. As far as the trend of average annual cosmic-ray intensity in Tbilisi is concerned, into 1963-1990 it took the form of the ninth power polynomial [43].

In table 2 is presented the statistical characteristics of real data of time-series of average annual air temperature, surface ozone concentration, galactic cosmic rays intensity and common mortality of the population of Tbilisi city.

As follows from table 2, the values of T varied from 12.4 up to 14.8 (average = 13.7), values of SOC – from 26.3 up to 61.8 (average = 44.1), values of CR – from 7803 up to 9100 (average = 8621) and M – from 8.03 up to 12.35 (average = 10.3).

The significant linear correlation M with SOC (positive) and CR (positive) are observed. Value of R between M and T is not significance.

Table 2

The statistical characteristics of real data of T, SOC, CR and M in Tbilisi in 1984-2010.

Variable	T	SOC	CR	M
Max	14.8	61.8	9100	12.35
Min	12.4	26.3	7803	8.03
Range	2.4	35.5	1297	4.32
Average	13.7	44.1	8621	10.3
St Dev	0.6	10.1	328	1.3
Correlation Matrix				
	T	SOC	CR	M
T	1	-0.08 (not sign)	0.02 (not sign)	0.01 (not sign)
SOC	-0.08 (not sign)	1	-0.26 ($\alpha=0.2$)	0.39 ($\alpha=0.05$)
CR	0.02 (not sign)	-0.26 ($\alpha=0.2$)	1	0.24 ($\alpha=0.2$)
M	0.01(not sign)	0.39 ($\alpha=0.05$)	0.24($\alpha=0.2$)	1

In table 3 are presented the statistical characteristics of the random components of T, SOC, CR and M in Tbilisi.

As follows from table 3, max and average values of random components of investigation parameters respectively are equal: T – 2.3 and 1.2, SOC - 14.7 and 7.3, CR – 587.5 and 314.9, M – 1.60 and 0.65.

The significant linear correlation only between M and T is observed (negative).

Table 3

The statistical characteristics of random components of T, SOC, CR and M in Tbilisi in 1984-2010.

Variable	T	SOC	CR	M
Max	2.3	14.7	587.5	1.60
Range	2.3	14.7	587.5	1.60
Average	1.2	7.3	314.9	0.65
St Dev	0.54	4.2	141	0.5
Correlation Matrix				
	T	SOC	CR	M
T	1	-0.30 ($\alpha=0.1$)	-0.40 ($\alpha=0.05$)	-0.41($\alpha=0.05$)
SOC	-0.30 ($\alpha=0.1$)	1	0.56 ($\alpha=0.005$)	0.11 (not sign)
CR	-0.40 ($\alpha=0.05$)	0.56 ($\alpha=0.005$)	1	0.13 (not sign)
M	-0.41($\alpha=0.05$)	0.11 (not sign)	0.13 (not sign)	1

Shares of the average values of random components from the average values of the real values of the investigated parameters (fig. 1-2, table 2 and 3) constitute: for T – 8.8%, for SOC – 16.6 %, for CR – 3.7 % and for M – 6.3 %.

The equation of the multiple linear regression of the connection of annual common mortality of the population of Tbilisi city with T_{rand} , SOC_{real} , and CR_{real} is represented below:

$$M = -0.3881 \cdot T_{rand} + 0.06074 \cdot SOC_{real} + 0.0013088 \cdot CR_{real} - 3.183$$

$$R^2 = 0.30 (\alpha=0.01).$$

In table 4 are presented data about contribution of variations in the values of T_{rand} , SOC_{real} , and CR_{real} to the changeability of M.

Table 4

Contribution of variations in the values of T_{rand} , SOC_{real} and CR_{real} to the changeability of M.

Variable	In the limits of Range (%)	In the limits of St Dev (%)
T_{rand}	8.5	4.0
SOC_{real}	20.9	11.9
CR_{real}	16.5	8.3

As follows from table 4, within the variation range the contribution of the studied parameters to annual common mortality of the population of Tbilisi city variability is as follows: random components of mean annual air temperature - 8.5%, real values of surface ozone concentration – 20.9%, real values of neutron component of galactic cosmic rays intensity - 16.5%. In the limits of standard deviation – 4.0 %, 11.9 % and 8.3 %, accordingly.

Conclusions

In the linear approximation the greatest contribution to the changeability of the annual mortality of population in Tbilisi city introduces the pollution of the atmosphere (ozone as direct contaminator and as the index of air quality). Significant role in these variations plays the changeability of the intensity of galactic cosmic rays (as the indicator of geomagnetic activity). The smallest contribution to the changeability of the mortality of the population of Tbilisi city introduce variations in the random component of the average annual air temperature.

References

- [1] Bunker A., Wildenhain J., Vandenberg A., Henschke N., Rocklöv J., Hajat Sh., Sauerborn R. Effects of Air Temperature on Climate-Sensitive Mortality and Morbidity Outcomes in the Elderly; a Systematic Review and Meta-analysis of Epidemiological Evidence. *EbioMedicine*, v. 6, 2016, pp. 258–268.
- [2] Turner, L.R., Barnett, A.G., Connell, D., Tong, S. Ambient temperature and cardio respiratory morbidity: a systematic review and meta-analysis. *Epidemiology*, v. 23, 2012, pp. 594–606.
- [3] Zhou X., Zhao A., Meng X., Chena R., Kuang X., Duan X., Kan H. Acute effects of diurnal temperature range on mortality in 8 Chinese cities. *Science of the Total Environment*, v. 493, 2014, pp. 92–97.
- [4] Wang X.Y., Barnett A.G., Hu W., Tong S. Temperature variation and emergency hospital admissions for stroke in Brisbane, Australia, 1996–2005. *Int. J. Biometeorol.*, v. 53, 2009, pp. 535–541. doi:10.1007/s00484-009-0241-4

- [5] Kyobutungi C., Grau A., Stieglbauer G., Becher H. Absolute temperature, temperature changes and stroke risk: a case-crossover study. *Eur. J. Epidemiol.*, v. 20, 2005, pp. 693–698. doi:10.1007/s10654-005-0703
- [6] Zaninović K., Matzarakis A. Impact of heat waves on mortality in Croatia. *Int. J. Biometeorol.*, v. 58, iss. 6, 2014, pp. 1135–1145.
- [7] Gomez-Acebo I., Llorca J., Dierssen T. Cold-related mortality due to cardiovascular diseases, respiratory diseases and cancer: a case-crossover study. *Public Health*, v. 127, 2013, pp. 252–258.
- [8] Amiranashvili A., Chikhladze V., Kartvelishvili L., Khazaradze K. Expected Change of the Extremal Air Temperature and its Influence on the Mortality (Based on the Example to Tbilisi City). International Cooperation Network for East European and Central Asian Countries: EECA Conference, Yerevan, Armenia, October 7-8, 2010, <http://be.sci.am/>.
- [9] Davis R.E., Knappenberger P.C., Novicoff W.M., Michaels P.J. Decadal changes in heat-related human mortality in the eastern United States. *Clim. Res.*, v. 22, 2002, pp. 175–184.
- [10] Huynen M.M.T.E., Martens P., Schram D., Weijenberg M.P., Kunst A.E. The Impact of Heat Waves and Cold Spells on Mortality Rates in the Dutch Population. *Environmental Health Perspectives*, v.109, N 5, 2001, pp. 463 – 470.
- [11] Davis R.E., Gregor G.R., Enfield K.B. Humidity: A review and primer on atmospheric moisture and human health. *Environmental Research*, v. 144, Part A, January 2016, pp. 106-116.
- [12] Amiranashvili A.G., Gogua R.A., Matiashvili T.G., Kirkitadze D.D., Nodia A.G., Khazaradze K.R., Kharchilava J.F., Khurodze T.V., Chikhladze V.A. The Estimation of the Risk of Some Astro-Meteo-Geophysical Factors for the Health of the Population of the City of Tbilisi. Int. Conference “Near-Earth Astronomy 2007” Abstract, Terskol, Russia, 3-7 September 2007, p. 86.
- [13] Shaposhnikov D., Revich B., Gurfinkel Yu., Naumova E. The influence of meteorological and geomagnetic factors on acute myocardial infarction and brain stroke in Moscow, Russia. *Int. J. of Biometeorology*, v. 58, iss. 5, 2014, pp. 799–808.
- [14] Azcárate T., Mendoza B., Levi J.R. Influence of geomagnetic activity and atmospheric pressure on human arterial pressure during the solar cycle 24. *Advances in Space Research*, v. 58, iss. 10, 2016, pp. 2116-2125.
- [15] Palmer S., Rycroft M., Cermack M. Solar and geomagnetic activity, extremely low frequency magnetic and electric fields and human health at the Earth’s surface. *Surv. Geophys.*, v. 27, 2006, pp. 557–595. doi:10.1007/s10712-006-9010-7
- [16] Caswell J.M., Carniello T.N., Murugan N.J. Annual incidence of mortality related to hypertensive disease in Canada and associations with heliophysical parameters. *Int. J. Biometeorol.*, v. 60, 2015, pp. 9–20. doi:10.1007/s00484-015-1000-3
- [17] Alexander L.T., Atawi N.S.A.I., Mostafa H.M.A. Space Weather Effects on Humans in Tabuk City, KSA. *Int. J. of Applied Science and Technology*, v. 6, N 1, February 2016, pp. 47-57.
- [18] Zenchenko T.A., Dimitrova S., Stoilova I., Breus T.K. Individual responses of arterial pressure to geomagnetic activity in practically healthy subjects. *Klin. Med.*, v. 87(4), 2009, pp.18–24.
- [19] Amiranashvili A., Bliadze T., Chikhladze V. Photochemical smog in Tbilisi. Monograph, Trans. of Mikheil Nodia institute of Geophysics, ISSN 1512-1135, v. 63, Tb., 2012, 160 p., (in Georgian).
- [20] Hori A., Hashizume M., Tsuda Y., Tsukahara T., Nomiya T. Effects of weather variability and air pollutants on emergency admissions for cardiovascular and cerebrovascular diseases. *Int. J. Environ Health Res.*, v. 22(5),2012, pp.416–430. doi:10.1080/09603123.2011.650155
- [21] Amiranashvili A., Matiashvili T., Nodia A., Nodia Kh., Kharchilava J., Khunjua A., Khurodze T., Chikhladze V. Air Electrical Conductivity Changeability as the Factor of Atmosphere Purity. Trans. of M. Nodia Institute of Geophysics, ISSN 1512-1135, v. 60, Tb., 2008, pp. 186 – 194, (in Russian).
- [22] Amiranashvili A., Chikhladze V., Bliadze T. Contemporary State of a Question About the Spatial-Temporary Distribution of Photochemical Smog and Ozone. Trans. of M. Nodia Institute of Geophysics, v. LXI, ISSN 1512-1135, Tb., 2009, pp. 179-186, (in Russian).

- [23] Amiranashvili A., Bliadze T., Melikadze G., Tarkhan-Mouravi I., Chikhladze V. Content of Light Aeroions as Factor of the Air Purity of Some Health Resorts of Georgia. *Modern Problems of Using of Health Resort Resources*, Collection of Scientific Works of International Conference, Sairme, Georgia, June 10-13, 2010, ISBN 978-9941-0-2529-7, Tbilisi, 2010, pp. 145-151, (in Russian).
- [24] Amiranashvili A., Chikhladze V., Bliadze T. Contemporary State of a Question About the Action of Photochemical Smog and Surface Ozone on Human Health. *Trans. of M. Nodia Institute of Geophysics*, v. LXII, ISSN 1512-1135, Tbilisi, 2010, pp. 177-188, (in Russian).
- [25] Amiranashvili A., Khurodze T., Shavishvili P., Beriashvili R. Iremashvili I. Dynamics of the Mortality of the Population of Tbilisi City and its Connection with the Surface Ozone Concentration. *Journ. of Georgian Geophysical Soc., Iss. (B), Physics of Atmosphere, Ocean and Space Plasma*, ISSN 1512-1127, v.16b, Tbilisi, 2013, pp. 31-38.
- [26] Amiranashvili A.G. Increasing Public Awareness of Different Types of Geophysical Catastrophes, Possibilities of Their Initiation as a Result of Terrorist Activity, Methods of Protection and Fight With Their Negative Consequences. *Engaging the Public to Fight Consequences of Terrorism and Disasters. NATO Science for Peace and Security Series E: Human and Societal Dynamics*, v. 120. IOS Press, Amsterdam•Berlin•Tokyo•Washington, DC, ISSN 1874-6276, 2015, pp.155-164. <http://www.nato.int/science>; <http://www.springer.com>; <http://www.iospress.nl>
- [27] Amiranashvili A., Amiranashvili V., Kartvelishvili L., Nodia Kh., Khurodze T. Influence of Air Effective Temperature and Geomagnetic Storms on the Population of Tbilisi City. *Trans. of the Institute of Hydrometeorology*, v. No 115, ISSN 1512-0902, Tbilisi, 2008, pp. 434 – 437, (in Russian).
- [28] Amiranashvili A., Mirianashvili K., Fedorova N., Levit V., Fabiana Medeiros Carnaúba, Aliton Oliveira da Silva. Comparative analysis of air equivalent - effective temperature in some cities of Georgia and Brazil, *Proc. of Int. Conf. “Environment and Global Warming”*, Dedicated to the 100th Birthday Anniversary of Academician F. Davitaya, *Collected Papers New Series*, N 3(82), ISSN 2333-3347, Tbilisi, 2011, pp. 105-110.
- [29] Amiranashvili A., Danelia R., Mirianashvili K., Nodia A., Khazaradze K., Khurodze T., Chikhladze V. On the Applicability of the Scale of Air Equivalent-Effective Temperature in the Conditions of Tbilisi City. *Trans. of M. Nodia Institute of Geophysics*, v. LXII, ISSN 1512-1135, Tbilisi, 2010, pp. 216-220, (in Russian).
- [30] Farajzadeh H., Saligheh M., Alijani B., Matzarakis A. Comparison of selected thermal indices in the northwest of Iran. *Natural Environment Change*, v. 1, N 1, 2015, pp. 1- 20.
- [31] Amiranashvili A.G., Chikhladze V.A. Saakashvili N.M., Tabidze M.Sh., Tarkhan-Mouravi I.D. Bioclimatic Characteristics of Recreational Zones – Important Component of the Passport of the Health Resort – Tourist Potential of Georgia. *Trans. of the Institute of Hydrometeorology at the Georgian Technical University*, v. 117, ISSN 1512-0902, Tbilisi, 2011, pp. 89-92.
- [32] Amiranashvili A.G., Bliadze T.G., Chikhladze V.A., Saakashvili N.M., Tarkhan-Mouravi I.D., Sikharulidze Sh.A., Lachashvili N.I. National Botanical Garden of Georgia – Recreational – Sanitation Oasis of Tbilisi City. *Trans. of the Institute of Hydrometeorology at the Georgian Technical University*, v. 117, ISSN 1512-0902, Tbilisi, 2011, pp. 94-96.
- [33] Matzarakis A., Rudel E., Zygmuntowski M., Koch E. Bioclimatic maps for tourism purposes. *Physics and Chemistry of the Earth*, v. 35, 2010, pp. 57–62.
- [34] Davis R.E., Knight D., Hondula D., Knappenberger P.C. A comparison of biometeorological comfort indices and human mortality during heat waves in the United States. Paper presented at Human biometeorology: The heat 17th conference on biometeorology and aerobiology, San Diego, CA, 2006, https://ams.confex.com/ams/BLTA/FBioA/techprogram/paper_110867.htm
- [35] Ketterer Ch., Matzarakis A. Human-biometeorological assessment of the urban heat island in a city with complex topography – The case of Stuttgart, Germany. *Urban Climate*, v. 10, Part 3, December 2014, pp. 573-584.

- [36] Urban A., Kyselý J. Comparison of UTCI with Other Thermal Indices in the Assessment of Heat and Cold Effects on Cardiovascular Mortality in the Czech Republic. *Int. J. Environ. Res. Public Health*, vol. 11, 2014, pp. 952-967.
- [37] Amiranashvili A.G., Cornélissen G., Amiranashvili V., Gheonjian L., Chikhladze V.A., Gogua R.A., Matiashvili T.G., Paatashvili T., Kopytenko Yu.A., Siegelova J., Dusek J., Halberg F. Circannual and circadecennian stages in mortality from cardiovascular causes in Tbilisi, Republic of Georgia (1980-1992). *Scriptamedica (Brno)*, 75, 2002, pp. 255-260.
- [38] Marti-Soler H., Gonseth S., Gubelmann C., Stringhini S., Bovet P., Chen P-Ch., Wojtyniak B., Paccaud F., Tsai D-H., Zdrojewski T., Marques-Vidal P. Seasonal variation of overall and cardiovascular mortality: A study in 19 countries from different geographic locations. *PLOS ONE*, v.9 (11), 2014, <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0113500>
- [39] Amiranashvili A., Chargazia Kh., Chikhladze V., Japaridze N., Khazaradze K. The monthly variations in mortality from the cardiovascular diseases in Tbilisi. *Georgian Medical News*, N 5 (242), 2015, pp. 53-59.
- [40] Kharchilava J., Chikhladze V., Chochishvili K., Chkhaidze G. Changeability of Surface Ozone Concentration in Tbilisi in 1984-2010. *Proc. Int. Sc. Conference on "Environment and Global Warming"*, Tbilisi, Georgia, 15-17 September 2011, Vakhushti Bagrationi Institute of Geography, Collected Papers, New Series, N 3(82), ISSN 2233-3347, Tbilisi, 2011, pp. 111-116.
- [41] Kobisheva N., Narovlianski G. Climatological processing of the meteorological information, Leningrad, Gidrometeoizdat, 1978, 294 p., (in Russian).
- [42] Kendall M.G. *Time-series*, Moscow, 1981, 200 p., (in Russian).
- [43] Amiranashvili A.G., Bakradze T.S., Ghlonti N.Ya., Khurodze T.V., Tuskia I.I. On the Connection of Annual Variations of the Intensity of Galactic Cosmic Rays with the Changeability of Cloudiness and Air Temperature in Tbilisi. *Journ. of Georgian Geophysical Soc., Iss. (B), Physics of Atmosphere, Ocean and Space Plasma*, ISSN 1512-1127, v.19b, Tbilisi, 2016, pp. 128-134.
- [44] Nurtaev B.S., Nurtaev L. Long Term Trends in Climate Variability of Caucasus Region. *Journ. of Georgian Geophysical Soc., Iss. (B), Physics of Atmosphere, Ocean and Space Plasma*, ISSN 1512-1127, v.19b, Tbilisi, 2016, pp. 79-89.
- [45] Budagashvili T., Karchava J., Gunia G., Intskirveli L., Kuchava T., Gurgenzidze M., Amiranashvili A., Chikhladze T. Inventory of Greenhouse Gas Emissions and Sinks. Georgia's Initial National Communication on Under the United Nations Framework Convection on Climate Change, Project GEO/96/G31, Tb., 1999, 137 p.
- [46] Amiranashvili A., Matcharashvili T., Chelidze T. Climate change in Georgia: Statistical and nonlinear dynamics predictions. *Journ. of Georgian Geophysical Soc., Iss.(A), Physics of Solid Earth*, v.15A, Tb., 2011-2012, pp. 67-87.
- [47] Amiranashvili A., Chikhladze V., Kartvelishvili L. Expected Change of Average Semi-Annual and Annual Values of Air Temperature and Precipitation in Tbilisi. *Journ. of Georgian Geophysical Soc., Iss. (B), Physics of Atmosphere, Ocean and Space Plasma*, ISSN 1512-1127, v. 13B, Tbilisi, 2009, pp. 50 – 54.
- [48] Amiranashvili A., Kartvelishvili L., Trofimenko L., Khurodze T. Statistical Structure of Mean Annual Air Temperature in Tbilisi and St.-Petersburg in 1850-2012. *Proc. of Int. Conf. "Modern Problems of Geography"*, Dedicated to the 80th Anniversary Since the Fondation of Vakhushti Bagrationi Institute of Geography, Collected Papers New Series, N 5(84), ISSN 2233-3347, Tbilisi, 2013, pp. 160-163.

**ჰაერის ტემპერატურის, მიწისპირა ოზონის კონცენტრაციის და
გალაქტიკური კოსმოსური სხივების ინტენსივობის საშუალოწლიური
მნიშვნელობების ვარიაციების ზემოქმედება მოსახლეობის
სიკვდილიანობაზე ქალაქ თბილისში**

ა.ამირანაშვილი, თ.ბაქრაძე, ნ. ბერიანიძე, ნ. ჯაფარიძე, ე. ხაზარაძე

რეზიუმე

წარმოდგენილია ჰაერის ტემპერატურის, მიწისპირა ოზონის კონცენტრაციის და გალაქტიკური კოსმოსური სხივების ნეიტრონული კომპონენტის ინტენსივობის საშუალოწლიური მნიშვნელობების ვარიაციების მოსახლეობის სიკვდილიანობაზე ზემოქმედების გამოკვლევის შედეგები ქალაქ თბილისში 1984–2010 წლებში. შესწავლილია გამოსაკვლევ მიწისპირა ოზონის სტატისტიკური მახასიათებლები. კერძოდ მიღებულია, რომ ვარიაციული განშლადობის ფარგლებში გამოსაკვლევ პარამეტრების წვლილი სიკვდილიანობის ცვალებადობაში შემდეგია: ჰაერის ტემპერატურის შემთხვევითი კომპონენტისა – 8.5%, მიწისპირა ოზონის კონცენტრაციისა და კოსმოსური სხივების ინტენსივობის რეალური მნიშვნელობებისა – 20.9% და 16.5% შესაბამისად.

**Влияние вариаций среднегодовых значений температуры
воздуха, концентрации приземного озона и интенсивности
галактических космических лучей на смертность населения
города Тбилиси**

**А.Г. Амиранашвили, Т.С. Бакрадзе, Н.Т. Берианидзе, Н.Д. Джапаридзе,
К.Р. Хазарадзе**

Резюме

Представлены результаты исследования влияния годовой изменчивости температуры воздуха, концентрации приземного озона и интенсивности нейтронной компоненты галактических космических лучей на смертность населения города Тбилиси в 1984-2010 гг. Изучены статистические характеристики исследуемых рядов. В частности, получено, что в пределах вариационного размаха вклад исследуемых параметров в изменчивость смертности следующий: случайной компоненты температуры воздуха – 8.5 %, реальных значений концентрации приземного озона и интенсивности космических лучей – 20.9 % и 16.5 % соответственно.